INDEX TO VOLUME 13

[New technical names in **bold-face** type]

A

Aceratherium gaudryi, 171.
Aepyceros malampus, 303.
Alces americana, 198.
americana americana, 29.
Alexander, Annie M., "Control," not "extermination," of Cynomys ludovicianus arizonensis, 302.
Allaceropinae, 171.
Allacerops, 170.
osborniana, 171.
turgaica, 171.
Allen, G. M., A pleistocene bat from Florida, 256. (See also Barbour, T.)
Alopex lagopus innuitus, 25.
Ammospermophilus interpres, 324, 326, 327.
anteater, 243.
antelope, 303.
pronghorn, 41–45.
sable, 303.
Antilocapra americana, 41.
Antrozous pallidus pallidus, 71.
Aotus, 368.
Aphelops, 265.
Apodemus sylvaticus, 218.
Archaeotherium, 358, 360.
Argyropulo, A. I., Preliminary description of two new palearctic voles, 268.
Arvicola, 151, 152, 280.
chrotorrhinus, 156.
Ashcraft, G. P., A third record of the spotted bat (Euderma maculata) for California, 162.
Ateles ater, 368.
dariensis, 367.
geoffroyi, 368.
grisescens, 367–368.

B

badger, 206, 329–330 (illus.).
Bailey, Florence M., Abert squirrel burying pine cones, 165.
Bailey, V., Trapping animals alive, 337.
Balaena cisarctica (type locality), 81–82.
Barbour, T., Concerning Ateles grisescens, 367.
Barbour, T., and G. M. Allen, The lesser one-horned rhinoceros, 144.
bat, 347–357, 363–365 (food).
big-eared, 71.
canyon, 325.
desert pallid, 71.
hoary (breeding habits), 365–367.
large brown, 71, 365.
large-eared free-tailed, 160.
little brown, 365.
Mexican bulldog, 159–160.
northern little brown, 202.
red (habits), 132–135.
spotted, 162–163.
bear, 329.
barren ground, 23.
black, 22, 202, 206, 274–275 (parasite).
grizzly, 22.
Pelly grizzly, 202.
beaver, 38, 201, 207, 117–126 (mental characteristics), 229–241 (fossil), 343–346 (illus., pond sites).
Canadian, 28.
Colorado River, 266–267.
Berry, L. J., A "horned" blacktail doe with fawn, 282.
bison, 40.
wood, 31, 198.

Bison bison athabascae, 31, 198.
Blarina brevicauda, 39.
brevicauda talpoides, 204.
costaricensis, 224.
pyrrhonota, 223, 225 (fig.).
thomasi, 224.
Blossom, P. M., A pair of long-tailed shrews (Sorex cinereus cinereus) in captivity, 136.
bobcat, 205.
Boochoerus, 358.
angustus, 359 (fig.), **360.**
humerosus, 360.
Brachitanytes, 369.
Brachylagus idahoensis, 79.
Bradt, G. W., The mammals of the Malpais, an area of black lava rock in the Tularosa basin, New Mexico, 321.
Bradypus tridactylus, 243.
buffalo, 186, 371–372 (disease).
African, 303.
wood, 31, 186.
Burt, W. H., The fish-eating habits of Pizonyx vivesi (Menegaux), 363; The systematic status and geographic range of the San Gabriel pocket gopher (Thomomys neglectus Bailey) 369.

C

Caenolestes fuliginosus, 226.
Cahalane, V. H., Large brown bat in Michigan, 70.
Cahn, A. R., Records and distribution of the fossil beaver, Castoroides ohioensis, 229.
Callospermophilus lateralis arizonensis, 220 (food), 222.
Canis latrans, 25, 205.
nubilus, 205.
occidentalis, 26, 201.
ochropus, 312.
caribou, 40.
barren ground, 30.
Osborn, 199.
Richardson's woodland, 31.
Carnivora, 53.
Castor canadensis, 38.
canadensis baileyi, 266.
canadensis canadensis, 28, 201, 207.
canadensis frondator, 266.
canadensis mexicanus, 266.
canadensis repentinus, 266–267.
Castoroides ohioensis (fig.), 229–241.
cattle, white park, 304.
Cervus canadensis canadensis, 199.
Chaerephon, 347.
Chaerodon, 358.
Cheiromeles, 347.
chickaree, 206.
chipmunk, antelope, 324, 325, 326, 327.
eastern, 166–167 (range), 278.
gray-collared, 222.
Lake Superior, 207.
Liard River, 27.
Lyster's, 207.
northern, 200
Chriacus (fig.), 53.
baldwini, 54.
Citellus grammurus tularosae, 326.
tridecemlineatus tridecemlineatus, 164–165, 207.
Clethrionomys, 280.
gapperi, 37, 281.
gapperi athabascae, 28, 200.
gapperi gapperi, 208, 209.
Cockerell, T. D. A., Kruger National Park, 303.
Colaptes cafer collaris, 221, 222.
comment and news, 97, 186, 302, 388.
Condylura cristata, 39, 204.
Connochaetes taurinus, 303.
Coolidge, H. J., Jr., Activities of the American committee for international wild life protection during the past year for the American Society of Mammalogists, 388.
Corynorhinus macrotis pallescens, 70.
Cosomys, 263.
glareolus, 281.
primus, 150–154, 280.

cougar, 72.
coyote, 205, 312, 329.
northern, 25.
Cryptotis, 224.
avia, 225 (fig.), 226.
equatoris, 225 (fig.), 226.
medellinus, 224, 225 (fig.), 226.
meridensis, 225 (fig.), 226.
merus, 224, 225 (fig.), 226, 227.
montivaga, 225 (fig.), 226, 227.
osgoodi, 224, 225 (fig.).
parva, 39.
squamipes, 224, 225 (fig.), 226.
thomasi, 225 (fig.), 226.
Cyanocitta stelleri diademata, 220, 222.
Cynomys ludovicianus arizonensis, 302.

D

Daeodon, 358.
calkinsi, 362.
minor (figs.), 360, 361-362.
shoshonensis, 362.
Dearborn, N., Occurrence of the least weasel in Michigan, 277.
deer, blacktail, 282-283.
fish-eating, 80.
mule, 29, 80 (birth date), 199.
white-tailed, 39.
Delphinus delphis, 69.
Diacodon septentrionalis, 51, 52 (fig.).
Dibelodon, 153.
Dice, L. R., The songs of mice, 187.
Diceratherium, 361.
Dinohyus, 358, 360, 361, 362.
Diphyllobothrium latum, 274-275.
Dipodomys deserti, 318.
ingens (habitat), 305-320.
merriami, 279.
merriami simiolus, 279.
nitratoides, 313.
ordii ordii, 324, 326.
spectabilis, 302, 305, 318.
spectabilis spectabilis, 309.
dogs, destructiveness, 98.
dolphin, 69.
Dysopes abrasus, 351.
gigas, 349.
glaucinus, 353.
perotis, 349.

E

Echidna, 242.
aculeata (anatomy, fig.), 55-58.
Ectypodus, 49, 51.
cochranensis (fig.), 49.
musculus, 51.
Eggysodon osborni, 170, 171.
pomeli, 170.
elephant, 303.
elk, 329, 331-336 (calls), 371-372 (disease).
Engels, W. L., Mid-winter activity in a striped spermophile, 164.
English, P. F., Some habits of the pocket gopher, Geomys breviceps breviceps, 126.
Engyodon, 170, 171.
entelodont (figs.), 358-362.
Eotrigonias, 170.
Epiaceratherium, 169-171.
bolcense, 170.
turgaicum, 170, 171.
Eptesicus fuscus, 365.
fuscus fuscus, 71, 161, 273.
Equus, 153.
burchelli transvaalensis, 303.
Erethizon dorsatum, 209.
epixanthum (song), 168-169.
epixanthum nigrescens, 201.
Euarctos americanus, 202, 206.
americanus americanus, 22, 274-275 (parasite).
Euderma maculata, 162-163.
maculatum, 163.
Eumops, 257, 347-357.
abrasus, 351.
abrasus abrasus, 349, 351-352.
abrasus milleri, 347 (breeding habits), 349, 352.
barbatus, 352.
bonariensis, 347, 354.

Eumops, **bonariensis beckeri, 349, 354–355, 356.**
bonariensis bonariensis, 349, 354, 355.
bonariensis delticus, 349, 355.
bonariensis nanus, 349, 356.
bonariensis patagonicus, 349, 355–356.
californicus, 274, 347, 351.
dabbenei, 350.
delticus, 355.
glaucinus, 349, 351, 352, 353–354.
hansae, 349, 356–357.
maurus, 348, 352–353.
milleri, 352.
nanus, 356.
orthotis, 353
patagonicus, 355.
perotis, 347, 349.
perotis californicus, 348, 351.
perotis dabbenei, 348, 350.
perotis perotis, 348, 349–350.
perotis trumbulli, 348, 350–351.
trumbulli, 350.
Eutamias asiaticus lineatus, 58.
asiaticus okadae, 58–59.
asiaticus orientalis, 59.
asiaticus uthensis, 59.
cinereicollis cinereicollis, 222.
minimus borealis, 27, 200.
minimus jacksoni, 207.
Eyerdam, W. J., A shipment of muskrats to Kamchatka, 281; Kamchatka sable nearing extinction, 276.

F

Felis concolor, 201.
fisher, 201.
flicker, red-shafted, 221.
flying squirrel, 206.
fox, British Columbia red, 24, 201.
Continental Arctic, 25.
gray, 205.
red, 205.
Fritz, E., Squirrel damage to young redwood trees, 76.

G

Geomys breviceps breviceps, 126–132 (habits), 128 (food), 130 (fig.).
bursarius illinoensis, 77–78.
pinetis, 78.
pinetis, 78.
Gianini, C. A., Birth date of mule deer fawns, 80.
gibbon, small black (nomenclature), 368–369.
Giraffa capensis, 303.
giraffe, 303.
Glaucomys sabrinus macrotis, 206.
Glyptotherium, 153.
Goldman, E. A., A new beaver from Arizona, 266; Review of wood rats of Neotoma lepida group, 59. (See also Nelson, E. W.)
Goodwin, G. G., New records and some observations on Connecticut mammals, 36.
Gordon, K., Rabbits killed on an Idaho highway, 169.
Gorsuch, D. M., 218.
Grange, W. B., Observations on the snowshoe hare, Lepus americanus phaeonotus Allen, 1; The pelage and color changes of the snowshoe hare, Lepus americanus phaeonotus Allen, 99.
Green, C. V., Breeding habits in captivity of Mus bactrianus Blyth, 45.
Green, M. M., The Franklin spermophile in Ontario, 277.
Gregory, T., In pursuit of badgers, 329.
Grinnell, J., Habitat relations of the giant kangaroo rat, 305.
ground squirrel, 329.
Arizona mantled, 220 (food), 222.
Douglas, 73–74, 212.
Gudger, E. W., 186.
Gulo luscus, 23, 202.

H

Hall, E. R., A new shrew of the Sorex merriami group from Arizona, 259.

hare, European, 38.
Hudson Bay varying, 29.
Keewatin Arctic, 29.
snowshoe, 1–19 (habits), 12 (illus.), 99–116 (pelage), 114 (figs.).
varying, 38, 209.
Harper, F., Mammals of the Athabaska and Great Slave Lakes region, 19.
Hesperomys cognatus, 192.
michiganensis, 192.
Hinton, M. A. C., Note on "Cosomys" Wilson, from the pliocene of California, 280.
hippopotamus, 303.
Hippotragus, 303.
Holt, E. G., Swimming cats, 72.
Howell, A. H., Notes on range of the eastern chipmunks in Ohio, Indiana and Quebec, 166.
Hudson, G. E., On the food habits of Marmosa, 159.
Huestis, R. R., Local differences in pelage characters of Peromyscus, 210.
Huey, L. M., Occurrence of Mexican bulldog bat in Panama, 159; The large-eared free-tailed bat in San Diego, 160.
Hyaenognathus dubius, 263.
matthewi, 263.
pachyodon, 263.
solus (fig.), 263–266.
Hylobates, 368.
Hylobatidae, 368.

I

impalla, 303.
Insectivora, 51.

J

Jacobi, A., Das Rentier (rev., Miller), 173.
jaguar, 72.
jay, long-crested, 220, 222.
Johnson, C. E., Notes on a family of red bats in captivity, 132.
junco, 222.
Junco phaeonotus dorsalis, 222.

K

Komarek, E. V., Distribution of Microtus chrotorrhinus, with description of a new subspecies, 155; Notes on mammals of Menominee Indian Reservation, Wisconsin, 203.
Kruger National Park, 303–304.
Kuroda, N., A description of an apparently new form of chipmunk from South Kuriles, Japan, 58.

L

Lagurus curtatus, 167.
curtatus artemisiae, 167.
intermedius, 167.
Lama, 153.
Langworthy, O. R., The panniculus carnosus and pouch musculature of the opossum, a marsupial, 241.
Lasiurus borealis borealis, 132–135 (habits), 162.
cinereus (breeding habits), 365–367.
seminola, 162.
seminolus, 162.
Leighton, A. H., Notes on the beaver's individuality and mental characteristics, 117.
lemming, tawny, 200.
Lemmus helvolus, 200.
Leo leo melanochaitus, 303.
Leptictidae, 51.
Lepus americanus, 38.
americanus americanus, 29, 199.
americanus phaeonotus, 1, 99–116 (pelage), 114 (figs.).
arcticus canus, 29.
lion, 303.
Loomis, F. B., Two new Miocene entelodonts, 358.
Lutra canadensis canadensis, 24, 205.
lynx, 205.
bay, 39.
Canada, 26, 39.
Lynx canadensis, 39.
canadensis canadensis, 26, 205.
rufus, 39, 205.

Lyon, M. W., Jr., Remarks on Geomys bursarius illinoensis Komarek and Spencer, 77.

M

Macropus bennetti, 245.
Mailliard, J., Ground squirrels invading new territory in the San Francisco Bay region, 73.
Marmosa zeledoni (food), 159.
marmot, Robson hoary, 200.
Marmota caligata oxytona, 200.
monax canadensis, 26.
monax rufescens, 207.
marten, Alaska pine, 201.
Martes americana actuosa, 201.
pennanti pennanti, 201.
mastodon, 263.
McKee, E. D., Big-eared bat in the Grand Canyon, 71.
Megachoerus, 358.
Mephitis hudsonica, 24, 206.
Merriam, C. Hart, 97 (letter).
Merycodus, 153.
Microtus, 37, 212, 280.
chrotorrhinus (distribution) 155–158.
chrotorrhinus carolinensis, 158.
chrotorrhinus chrotorrhinus, 156–157.
chrotorrhinus ravus, 157.
colchicus, 268–271, 269 (fig.), 270 (fig.).
eversmanni, 268.
gregalis angustus, 268.
gregalis brevicauda, 268.
gregalis buturlini, 268.
gregalis castaneus, 268.
gregalis eversmanni, 268.
gregalis gregalis, 268.
gregalis kossogolicus, 268.
gregalis montosus, 268.
gregalis raddei, 268.
gregalis ravidulus, 268.
gregalis slowzovi, 268.
gregalis tianschanicus, 268.
gregalis unguiculatus, 268.
Microtus hartingi, 271.
lydius, 271.
major, 268.
mordax mordax, 201.
mustersi, 271.
pennsylvanicus drummondii, 200.
pennsylvanicus pennsylvanicus, 208.
philistinus, 271.
socialis guentheri, 271.
socialis irani, 271.
socialis paradoxus, 271.
socialis satunini, 271.
Miller, G. S., Jr., Das Rentier (Jacobi, reviewed), 173; The search for man's ancestors (Smith, reviewed), 373; The social life of monkeys and apes (Zuckerman, reviewed), 83.
Mimomys, 151, 280.
intermedius, 280.
pliocaenicus, 281.
pliocoenicus, 152.
primus, 281.
Mimon, 257.
mink, 206.
northwestern, 201.
Preble's, 23.
Mohr, C. E., Myotis grisescens and Myotis sodalis in Tennessee and Alabama, 271; Myotis subulatus leibii and Myotis sodalis in Pennsylvania, 160–161.
mole, Brewer's, 38.
common garden, 38.
star-nosed, 39, 204.
Molossidae, 347.
Molossides, 257.
floridanus (fig.), **257–259.**
Molossops temmincki, 258.
Molossus, 257, 347.
abrasus, 351.
bonariensis, 354.
californicus, 348, 351.
ferox, 353.
gigas, 349.
glaucinus, 353.
maurus, 352.

Molossus obscurus, 258.
perotis, 349.
monkey, Balboa, 368.
gray spider, 367–368.
howling (parasite), 368.
night, 368.
North American black spider, 367.
red spider, 368.
Monodelphis, 223.
moose, 40, 198, 329.
eastern, 29.
Mormopterus, 347.
mountain goat, 198.
lion, 201.
sheep, Stone, 197–198.
mouse, Athabaska red-backed, 28.
Baird pocket, 324, 325, 327.
black Malpais pocket, 327.
boreal white-footed, 28.
cactus, 324, 326.
Canadian deer, 208.
cantankerous meadow, 201.
Cooper's lemming, 209.
deer, 192, 222.
desert pocket, 325.
Drummond meadow, 200.
golden harvest, 192.
harvest, 192.
house, 187–196 (song), 208.
Hudson Bay jumping, 208.
jumping, 201.
long-nosed deer, 324, 326, 327.
Malpais pocket, 324, 325.
northern meadow jumping, 29.
northern white-footed, 200, 208.
pinyon, 324, 326, 327.
pocket, 279, 321.
red-backed, 200, 208.
rock pocket, 324, 325, 326, 327.
San Martin Island (nomenclature), 370–371.
short-tailed meadow, 167.
white-footed, 222.
wood, 218.
Multituberculata, 49.
Murie, O. J., Elk calls, 331; and A. Murie, Further notes on travels of Peromyscus, 78
Murie, A., 78 (see also Murie, O. J.)
Murray, L. T., Notes on personal experiences with pronghorn antelope in Texas, 41.
Mus bactrianus (habits), 45–47.
gansuensis, 45.
musculus, 45, 187–196 (song), 208.
tantillus, 45.
musk-ox, 35.
muskrat, 208, 281–282.
northwestern, 28.
Mustela arizonensis, 71.
cicognanii, 201.
cicognanii richardsonii, 23.
noveboracensis, 163–164.
rixosa allegheniensis, 277.
vison, 206.
vison energumenos, 201.
vison lacustris, 23.
Myopterus maurus, 352.
Myotis grisescens, 272–273.
incautus, 70.
keenii septentrionalis, 202
lucifugus, 365.
lucifugus lucifugus, 161.
sodalis, 160–161, 272–273.
subulatus leibii, 160–161.
thysanodes, 70.

N

Napaeozapus insignis, 36, 37.
Nelson, E. W. and E. A. Goldman, The name of the San Martin Island mouse —a correction, 370; The type locality of Ursus lotor Linne, 367.
Neofiber, 152.
Neosorex palustris navigator, 203.
Neotoma abbreviata, 65.
albigula albigula, 326.
albigula melas, 321, 324, 326, 327.
anthonyi, 66.
arenacea, 65.
bella, 62.
bella felipensis, 64.
bryanti, 65.
californica, 64.
cinerea drummondi, 200.
desertorum, 61, 62.

Neotoma desertorum sola, 63.
goldmani, 67.
insularis, 65.
intermedia, 64.
intermedia devia, 62.
intermedia gilva, 63.
intermedia notia, 65.
intermedia perpallida, 65.
intermedia pretiosa, 64.
intermedia ravida, 64.
intermedia vicina, 65.
lepida, 59, 61.
lepida arenacea, 65.
lepida devia, 62.
lepida felipensis, 64.
lepida gilva, 63.
lepida intermedia, 64.
lepida lepida, 61, 64.
lepida monstrabilis, 62–63.
lepida notia, 65.
lepida perpallida, 65.
lepida pretiosa, 64.
lepida ravida, 64.
lepida vicina, 65.
martinensis, 66.
micropus canescens, 70, 324, 326.
nevadensis, 62.
nudicauda, 65.
pennsylvanica, 38.
stephensi, 66.
stephensi relicta, 66–67.
stephensi stephensi, 66.
Noctilio, 257, 363.
leporinus mexicanus, 159–160
Nyctinomus depressus, 160.
orthotis, 353.

O

Ocelot, 72.
Odocoileus, 153.
columbianus, 282–283.
hemionus, 199.
hemionus hemionus, 29.
virginianus, 39.
Oligoryzomys, 226.
Olson, S. F., Fish-eating deer, 80.
Ondatra zibethica spatulata, 28.
Ondatra zibethica zibethica, 208.
Onychomys torridus ramona, 78.
opossum (anatomy, figs.), 241–250.
Orcinus rectipinna, 171–172.
Oreamnos americanus columbiae, 198.
Ornithorhynchus, 242.
Orr, R. T., Occurrence of the short-tailed meadow mouse in northern California, 167.
Otospermophilus grammurus, 326.
grammurus douglasii, 73–74.
grammurus tularosae, 324–327.
otter, 205.
Canadian, 24.
Ovibos moschatus moschatus, 35.
Ovis stonei, 197–198.
Oxyclaenidae (fig.), 53.
Oxyclaenus, 53.

P

Palmer, T. S., 388, Obituary notice of Hermann von Wechlinger Schulte.
Parascalops breweri, 36, 38.
Parectypodus simpsoni, 51.
Parker, H. C., Notes on mammals of the Carlsbad Cavern region, 70.
Parker, G. H., The respiratory rate of the common porpoise, 68.
peccary, 72, 263.
Pelonax, 358.
Perodipus ingens, 305.
Perognathus, 279.
baileyi baileyi, 279–280.
californicus, 279.
fallax, 279.
flavus, 325, 327.
flavus flavus, 324.
formosus, 279.
formosus mesembrinus, 279.
intermedius, 327.
intermedius ater, 321, 324, 325, 327.
intermedius intermedius, 324.
longimembris, 279.

Perognathus longimembris arenicola, 279.
penicillatus, 279.
penicillatus angustirostris, 279.
penicillatus eremicus, 325.
spinatus, 279.
spinatus rufescens, 279.
Peromyscus, 210–218 (pelage), 213 (fig.), 215 (fig.), 219.
crinitus stephensi, 210, 279.
eremicus, 215, 279, 326.
eremicus eremicus, 279, 324.
exiguus, 371.
gossypinus, 192.
leucopus noveboracensis, 208.
leucopus tornillo, 70.
maniculatus artemisiae, 78.
maniculatus bairdii, 192.
maniculatus borealis, 28, 200.
maniculatus exiguus (nomenclature), 371.
maniculatus gambeli, 216.
maniculatus geronimensis, 371.
maniculatus gracilis, 208.
maniculatus martinensis (nomenclature), 370–371.
maniculatus rufinus, 210, 222.
nasutus, 326, 327.
nasutus griseus, 324, 326.
polionotus albifrons, 211.
polionotus polionotus, 211.
truei, 326, 327.
truei truei, 324.
Phacochoerus aethiopicus sundevalli, 303.
Phaiomys, 281.
Phocaena phocoena (respiration), 68–69.
Pipistrellus hesperus hesperus, 273 (roost), 325.
subflavus, 161.
subflavus subflavus, 161, 273.
Pitymys, 280.
Pizonyx vivesi, 273, 363–365 (food).
Platymops, 347.
Plesippus, 263.
Pliauchenia, 153.
Pliohippus, 263.
pocket gopher, 126–132 (habits), 128 (food), 130 (figs.).
San Gabriel (range), 369–370.
Poole, E. L., Breeding of the hoary bat in Pennsylvania, 365; Lasiurus seminolus in Pennsylvania, 162.
porcupine, 38, 167–168 (voice), 168–169 (song).
Canada, 209.
dusky, 201.
porpoise, bottle-nosed, 68.
common (respiration), 68–69.
prairie dog, 74.
white, 75.
Wilcox, 302.
Preptoceras, 265.
Procamelus, 153.
Procyon lotor, 367.
lotor lotor, 205.
Prohyracodon, 170.
Promops, 347.
abrasus, 351.
barbatus, 352.
bonariensis, 354.
californicus, 351.
glaucinus, 353.
milleri, 352.
nanus, 356.
orthotis, 353.
perotis, 349.
trumbulli, 350.
Ptilodontidae, 49.
Ptilodus (fig.), 49.
cochranensis, 49.
gracilis, 51.
trovessartianus, 51.
puma, 39, 72.

R

rabbit, 221.
desert cottontail, 325.
Idaho pygmy, 79.
jack, 169, 329.
Mearns' cottontail, 209.
New England cottontail, 38.
snowshoe, 199.
raccoon, 205, 367.
Radcliffe, L., Status of international convention for protection of whales, 251.
Rangifer arcticus arcticus, 30.
caribou sylvestris, 31.
osborni, 199.
rat, giant kangaroo (habitat), 305–320.
kangaroo, 309.
Norway, 208.
Ord kangaroo, 324, 326.
rats transporting eggs, 186.
Rattus norvegicus, 46, 208.
recent literature, 83, 173, 290, 373.
Reithrodontomys fulvescens aurantius, 192.
megalotis, 192.
rhinoceros, 169.
black, 303.
Indian, 144.
lesser one-horned, 144–149, 148 (fig.).
two-horned, 145.
Rhinoceros sondaicus, 144–149, 148 (fig.).
sumatrensis, 145.
unicornis, 144.
Rhinocerotidae, 169.
Ronzotherium gaudryi, 170.
Rush, W. M., Bang's disease in the Yellowstone National Park buffalo and elk herds, 371; Diphyllobothrium latum in bear, 274.
Russell, L. S., New data on the paleocene mammals of Alberta, Canada, 48.

S

sable, 276–277.
Sanborn, C. C., The bats of the genus Eumops, 347.
Saunders, A. A., The voice of the porcupine, 167.
Scalopus aquaticus, 38.
Schultz, A. H., The generic position of Symphalangus klossii, 368.
Sciurus carolinensis leucotis, 206.
fremonti fremonti (food), 278–279.
griseus griseus, 76.
hudsonicus hudsonicus, 27, 199.
hudsonicus minnesota, 206.
niger, 38.
niger rufiventer, 206.
Seton, E. T., The song of the porcupine (Erethizon epixanthum), 168.
Sheldon, W. G., Mammals collected or observed in the vicinity of Laurier Pass, B. C., 196.
Shepherd, G. S., Killer whale in slough at Portland, Oregon, 171.
shrew, common, 204.
dusky, 202.
long-tailed (habits), 136–143.
short-tailed, 204.
water, 203.
skunk, Hudsonian, 206.
Northern Plains, 24.
sloth, three-toed, 243.
Smith, G. E., The search for man's ancestors (rev., Miller), 373.
Sorex brasiliensis, 223.
cinereus, 39.
cinereus cinereus, 136–143 (habits), 204.
dispar, 39.
fumeus, 36, 39.
leucogenys, 260.
melanogenys (fig.), **260–262.**
merriami, 259.
obscurus obscurus, 202.
palustris, 36, 39.
surinamensis, 223.
spermophile, Franklin, 277–278.
striped, 164–165.
thirteen-lined, 207.
squirrel, 221.
Abert (cache), 165–166.
dark rock, 327.

squirrel, fox, 38.
Fremont's (food), 278–279.
Hudson Bay red, 27.
northern gray, 206.
northern red, 199.
red, 206.
rock, 325, 326, 327.
western fox, 206.
Stanford, J. S., Idaho pygmy rabbit: Brachylagus idahoensis (Merriam), 79.
Stenocranius eversmanni, 268.
gregalis montosus, 268.
major, 268.
Stenomylus hitchcocki, 361.
Stephl, O. E., Battle between prairie dog and rattlesnake, 74; White prairie dogs found in Montana, 75.
Stock, C., Hyaenognathus from the late pliocene of the Coso Mountains, California, 263.
Stone, W., The type locality of Balaena cisarctica Cope, 81.
Sylvilagus auduboni minor, 325.
floridanus mallurus, 38.
floridanus mearnsi, 209.
mallurus, 36.
transitionalis, 38.
Symphalangus klossii (nomenclature), 368–369.
Synaptomys cooperi, 36.
cooperi cooperi, 209.

T

Tadarida, 347.
macrotis, 160.
orthotis, 353.
Tamandua tetradactyla, 243.
Tamias striatus, 278.
striatus fisheri (range), 166.
striatus griseus (range), 167.
striatus lysteri, 207.
Tate, G. H. H., Distribution of the South American shrews, 223.
Taxidea taxus, 206.
Taylor, W. P., and D. M. Gorsuch, A test of some rodent and bird influences on western yellow pine reproduction at Fort Valley, Flagstaff, Arizona, 218.
Test, F. H., Winter activities of the eastern chipmunk, 278.
Thomasomys baeops, 226.
Thomomys bottae bottae, 369.
bottae mohavensis, 369.
bottae neglectus, 370.
bottae pallescens, 76, 369.
bottae pascalis, 369.
mearnsi, 302.
neglectus (range), 369–370.
perpallidus mohavensis, 77.
perpallidus perpes, 77.
traps, 337–342.
Tricentes, 53.
Trigonias ombonii, 169.
Tursiops truncatus, 68, 69.

U

Urocyon cinereoargenteus, 205.
Ursus, 23.
lotor (type locality), 367.
pellyensis, 202.
richardsoni, 23.

V

vole, 150–154, 263.
gray rock, 157.
meadow, 208.
rock, 156–157.
Smoky Mountain rock, 158.
von Bloeker, J. C., Jr., Extension of the ranges of three pocket gophers in southern California, 76; New locality records for Perognathus baileyi from California, 279; Northern extension of the range of Onychomys torridus ramona Rhoads, 78; Summer records of Eumops californicus (Merriam), 274; The roosting-place of Pipistrellus hesperus, 273; Unusual abundance of pocket mice at San Felipe Narrows, California, 279.
Vulpes alascensis abietorum, 24, 201.
fulva fulva, 205.

W

wapiti, 199.
Warren, E. R., A somewhat unusual kitchen midden of Fremont's squirrel (Sciurus fremonti fremonti), 278; The abandonment and reoccupation of pond sites by beavers, 343; When do weasels mate? 71.
wart hog, 303.
weasel (attack of), 163–164.
    Bonaparte, 201.
    least, 277.
    mountain, 71.
    Richardson's, 23.
whales, 251–255.
Wight, H. M., A weasel attacks a man, 163.
wildebeest, 303.
Willis, R. A., A left-sided inferior vena cava in Echidna aculeata, 55.
Wilson, R. W., Cosomys, a new genus of vole from the Pliocene of California, 150.
wolf, 39, 205.
wolf, gray, 329.
    northern gray, 26, 201.
    northern timber, 26.
wolverene, Hudson Bay, 23.
wolverine, 202.
whale, killer, 171–172.
Wood, H. E., 2nd, Status of Epiaceratherium (Rhinocerotidae), 169.
woodchuck, 207.
        Canadian, 26.
wood rat, 38, 321.
    black, 325, 327.
    Drummond, 200.
    hoary, 325, 326.
    Malpais, 324.
    white-throated, 326, 327.

Z

Zapus hudsonius, 37.
    hudsonius hudsonius, 29, 208.
    princeps, 201.
zebra, 303.
Zuckerman, S., The social life of monkeys and apes (rev., Miller), 83.